

Neutrino Mass Matrix from S_4 Symmetry

Ernest Ma

*Physics Department, University of California, Riverside, California 92521, USA,
and Institute for Particle Physics Phenomenology, Department of Physics,
University of Durham, Durham, DH1 3LE, UK*

Abstract

The cubic symmetry S_4 contains A_4 and S_3 , both of which have been used to study neutrino mass matrices. Using S_4 as the family symmetry of a complete supersymmetric theory of leptons, it is shown how the requirement of breaking S_4 at the seesaw scale without breaking supersymmetry enforces a special form of the neutrino mass matrix which exhibits maximal $\nu_\mu - \nu_\tau$ mixing as well as zero U_{e3} . In addition, $(\nu_e + \nu_\mu + \nu_\tau)/\sqrt{3}$ is naturally close to being a mass eigenstate, thus predicting $\tan^2 \theta_{12}$ to be near but not equal to 1/2.

In atmospheric neutrino oscillation data [1], the persistence of maximal $\nu_\mu - \nu_\tau$ mixing has raised the important theoretical question of whether it is due to an underlying symmetry. The naive response is that it is due to the exchange of ν_μ with ν_τ , but since ν_μ and ν_τ are respective members of the $SU(2)_L$ doublets (ν_μ, μ) and (ν_τ, τ) , this symmetry automatically implies the exchange of μ with τ . As such, it cannot be sustained in the full Lagrangian, because the assumed diagonality of the charged-lepton mass matrix (with $m_\mu \neq m_\tau$) is then lost.

A related issue is whether or not $U_{e3} = 0$. If $\nu_\mu - \nu_\tau$ exchange were a legitimate symmetry, then this would also be “predicted”. On the other hand, $U_{e3} = 0$ by itself can be supported by a different symmetry of the full Lagrangian [2], although the latter is generally unable to shed any light on how maximal $\nu_\mu - \nu_\tau$ mixing could be achieved.

With the implementation of non-Abelian discrete symmetries such as S_3 [3], D_4 [4], and A_4 [5], as well as Q_4 [6] and Q_6 [7], it is indeed possible to have both maximal $\nu_\mu - \nu_\tau$ mixing and zero U_{e3} , as well as a prediction of the mixing angle in solar neutrino oscillations [8] in some cases. However, specific *ad hoc* assumptions regarding the symmetry breaking sector must be made, usually with the addition of arbitrary auxiliary Abelian discrete symmetries [9].

In this paper, the problem is solved by invoking a very simple requirement. The complete theory of leptons is assumed to be supersymmetric with S_4 as its family symmetry. Neutrino masses are assumed to come from the canonical seesaw mechanism [10] with heavy singlet neutral fermions N . The key is to allow S_4 to be broken by the Majorana mass matrix of N at the seesaw scale, but not the supersymmetry. This requirement then fixes the pattern of symmetry breaking and thus the form of $\mathcal{M}_N$, and subsequently also $\mathcal{M}_\nu$, as shown below.

The group of the permutation of four objects is S_4 [11]. It is also the symmetry group of the hexahedron, i.e. the cube, one of five (and only five) perfect geometric solids, which

was identified by Plato with the element “earth” [12]. It has 24 elements divided into 5 equivalence classes, with $\underline{1}$, $\underline{1}'$, $\underline{2}$, $\underline{3}$, and $\underline{3}'$ as its 5 irreducible representations. Its character table is given below.

Table 1: Character table of S_4 .

Class	n	h	χ_1	$\chi_{1'}$	χ_2	χ_3	$\chi_{3'}$
C_1	1	1	1	1	2	3	3
C_2	3	2	1	1	2	-1	-1
C_3	8	3	1	1	-1	0	0
C_4	6	4	1	-1	0	-1	1
C_5	6	2	1	-1	0	1	-1

The two three-dimensional representations differ only in the signs of their C_4 and C_5 matrices. Their group multiplication rules are similar to those of A_4 [13], namely

$$\underline{3} \times \underline{3} = \underline{1} + \underline{2} + \underline{3}_S + \underline{3}'_A, \quad (1)$$

$$\underline{3}' \times \underline{3}' = \underline{1} + \underline{2} + \underline{3}_S + \underline{3}'_A, \quad (2)$$

$$\underline{3} \times \underline{3}' = \underline{1}' + \underline{2} + \underline{3}'_S + \underline{3}_A, \quad (3)$$

where the subscripts S and A refer to their symmetric and antisymmetric product combinations respectively. The two-dimensional representation behaves exactly as its S_3 counterpart, namely

$$\underline{2} \times \underline{2} = \underline{1} + \underline{1}' + \underline{2}. \quad (4)$$

The three N 's are assigned to the $\underline{3}$ representation. To obtain a nontrivial $\mathcal{M}_N$, the singlet Higgs superfields $\sigma_{1,2,3} \sim \underline{3}$ and $\zeta_{1,2} \sim \underline{2}$ are assumed. Consequently in the $N_{1,2,3}$ basis,

$$\mathcal{M}_N = \begin{pmatrix} A + f(\langle \zeta_2 \rangle + \langle \zeta_1 \rangle) & h\langle \sigma_3 \rangle & h\langle \sigma_2 \rangle \\ h\langle \sigma_3 \rangle & A + f(\omega\langle \zeta_2 \rangle + \omega^2\langle \zeta_1 \rangle) & h\langle \sigma_1 \rangle \\ h\langle \sigma_2 \rangle & h\langle \sigma_1 \rangle & A + f(\omega^2\langle \zeta_2 \rangle + \omega\langle \zeta_1 \rangle) \end{pmatrix}, \quad (5)$$

where $\omega = \exp(2\pi i/3)$. The most general S_4 -invariant superpotential of σ and ζ is given by

$$\begin{aligned} W = & \frac{1}{2}M(\sigma_1\sigma_1 + \sigma_2\sigma_2 + \sigma_3\sigma_3) + \lambda\sigma_1\sigma_2\sigma_3 + m\zeta_1\zeta_2 + \frac{1}{3}\rho(\zeta_1\zeta_1\zeta_1 + \zeta_2\zeta_2\zeta_2) \\ & + \frac{1}{2}\kappa(\sigma_1\sigma_1 + \omega\sigma_2\sigma_2 + \omega^2\sigma_3\sigma_3)\zeta_2 + \frac{1}{2}\kappa(\sigma_1\sigma_1 + \omega^2\sigma_2\sigma_2 + \omega\sigma_3\sigma_3)\zeta_1. \end{aligned} \quad (6)$$

The resulting scalar potential is then

$$\begin{aligned} V = & |M\sigma_1 + \lambda\sigma_2\sigma_3 + \kappa\sigma_1(\zeta_2 + \zeta_1)|^2 \\ & + |M\sigma_2 + \lambda\sigma_3\sigma_1 + \kappa\sigma_2(\omega\zeta_2 + \omega^2\zeta_1)|^2 \\ & + |M\sigma_3 + \lambda\sigma_1\sigma_2 + \kappa\sigma_3(\omega^2\zeta_2 + \omega\zeta_1)|^2 \\ & + |m\zeta_1 + \rho\zeta_2\zeta_2 + \frac{1}{2}\kappa(\sigma_1\sigma_1 + \omega\sigma_2\sigma_2 + \omega^2\sigma_3\sigma_3)|^2 \\ & + |m\zeta_2 + \rho\zeta_1\zeta_1 + \frac{1}{2}\kappa(\sigma_1\sigma_1 + \omega^2\sigma_2\sigma_2 + \omega\sigma_3\sigma_3)|^2. \end{aligned} \quad (7)$$

For supersymmetry to be unbroken, $V_{min} = 0$ is required. This is possible only if $\langle\zeta_1\rangle = \langle\zeta_2\rangle$ and $\langle\sigma_2\rangle = \langle\sigma_3\rangle$, for which

$$M\langle\sigma_1\rangle + \lambda\langle\sigma_2\rangle^2 + 2\kappa\langle\sigma_1\rangle\langle\zeta_1\rangle = 0, \quad (8)$$

$$M + \lambda\langle\sigma_1\rangle - \kappa\langle\zeta_1\rangle = 0, \quad (9)$$

$$m\langle\zeta_1\rangle + \rho\langle\zeta_1\rangle^2 + \frac{1}{2}\kappa(\langle\sigma_1\rangle^2 - \langle\sigma_2\rangle^2) = 0, \quad (10)$$

where $\omega + \omega^2 = -1$ has been used. Thus $\mathcal{M}_N$ is fixed to be of the form

$$\mathcal{M}_N = \begin{pmatrix} A + 2B & C & C \\ C & A - B & D \\ C & D & A - B \end{pmatrix}, \quad (11)$$

where $B = f\langle\zeta_1\rangle$, $C = h\langle\sigma_2\rangle$, and $D = h\langle\sigma_1\rangle$.

The residual symmetry of the theory is then Z_2 , under which $(N_2 - N_3)$, $(\sigma_2 - \sigma_3)$, $(\zeta_1 - \zeta_2)$ are odd, and N_1 , $(N_2 + N_3)$, σ_1 , $(\sigma_2 + \sigma_3)$, $(\zeta_1 + \zeta_2)$ are even. If $\langle\zeta_{1,2}\rangle = 0$, then $B = 0$, $C = D$, and the residual symmetry is S_3 , under which $(N_1 + N_2 + N_3)$, $(\sigma_1 + \sigma_2 + \sigma_3) \sim \mathbb{1}$;

$(N_1 + \omega N_2 + \omega^2 N_3, N_1 + \omega^2 N_2 + \omega N_3), (\sigma_1 + \omega \sigma_2 + \omega^2 \sigma_3, \sigma_1 + \omega^2 \sigma_2 + \omega \sigma_3), (\zeta_1, \zeta_2) \sim \underline{2}$. The Z_2 (and the approximate S_3) symmetry of $\mathcal{M}_N$ is preserved in $\mathcal{M}_N^{-1}$.

Consider now the leptons (ν_i, l_i) and l_j^c of the Standard Model. They are also assigned to the $\underline{3}$ representation of S_4 . As for the Higgs doublet superfields, they are assumed to be $\underline{1} + \underline{2}$, with one set coupling (ν_i, l_i) to l_j^c and the other coupling (ν_i, l_i) to N_j , as in any supersymmetric extension of the Standard Model. Nonzero vacuum expectation values of the scalar components of these Higgs superfields break the electroweak symmetry, but because of their S_4 structure, the $l_i l_j^c$ mass matrix is diagonal, as well as that of $\nu_i N_j$. In particular,

$$\begin{pmatrix} m_e \\ m_\mu \\ m_\tau \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} y_1 v_1 \\ y_2 v_2 \\ y_2 v_3 \end{pmatrix}, \quad (12)$$

where v_1 is the vacuum expectation value of the $\underline{1}$ representation, $v_{2,3}$ are those of the $\underline{2}$, and $y_{1,2}$ are their respective Yukawa couplings. In the νN sector, the vacuum expectation values of the corresponding Higgs $\underline{2}$ representation are assumed to be zero, thus $m_1 = m_2 = m_3 (= m_D)$ for all the Dirac neutrino masses. Using the seesaw mechanism, the observed Majorana neutrino mass matrix in the $\nu_{e,\mu,\tau}$ basis is then given by

$$\mathcal{M}_\nu = -\mathcal{M}_D \mathcal{M}_N^{-1} \mathcal{M}_D^T = \begin{pmatrix} a+2b & c & c \\ c & a-b & d \\ c & d & a-b \end{pmatrix}, \quad (13)$$

where

$$a = [-A^2 + B^2 + (2C^2 + D^2)/3] m_D^2 / \det \mathcal{M}_N, \quad (14)$$

$$b = [B(A - B) - (C^2 - D^2)/3] m_D^2 / \det \mathcal{M}_N, \quad (15)$$

$$c = C(A - B - D) m_D^2 / \det \mathcal{M}_N, \quad (16)$$

$$d = [D(A + 2B) - C^2] m_D^2 / \det \mathcal{M}_N, \quad (17)$$

$$\det \mathcal{M}_N = (A - B - D)[(A + 2B)(A - B + D) - 2C^2]. \quad (18)$$

This form of $\mathcal{M}_\nu$ is precisely that advocated in Ref. [14] on purely phenomenological grounds. As expected, the S_3 limit is obtained if $B = 0$ and $C = D$, for which $b = 0$ and $c = d$, resulting in $(\nu_e + \nu_\mu + \nu_\tau)/\sqrt{3}$ as a mass eigenstate. Note that S_4 as well as the residual Z_2 symmetry are assumed broken softly at the scale of supersymmetry breaking. This allows the electroweak vacuum expectation values to be chosen as they are.

In the basis $[\nu_e, (\nu_\mu + \nu_\tau)/\sqrt{2}, (-\nu_\mu + \nu_\tau)/\sqrt{2}]$, the neutrino mass matrix of Eq. (13) becomes

$$\mathcal{M}_\nu = \begin{pmatrix} a + 2b & \sqrt{2}c & 0 \\ \sqrt{2}c & a - b + d & 0 \\ 0 & 0 & a - b - d \end{pmatrix}, \quad (19)$$

which exhibits maximal $\nu_\mu - \nu_\tau$ mixing and zero U_{e3} as advertised. The mixing angle of the 2×2 submatrix (for all parameters real) can be simply read off as

$$\tan 2\theta_{12} = \frac{2\sqrt{2}c}{d - 3b} = \frac{2\sqrt{2}c}{c - 3b - (c - d)}, \quad (20)$$

which reduces to $2\sqrt{2}$ in the S_3 limit of $b = 0$ and $c = d$. This would imply $\tan^2 \theta_{12} = 1/2$, resulting in a mixing pattern proposed some time ago [15]. However in this limit, Δm_{atm}^2 vanishes as well [16]. Therefore, $\tan^2 \theta_{12}$ should not be equal to $1/2$, but since it is natural for b and $c - d$ to be small compared to c , its deviation from $1/2$ is expected to be small. For example,

$$(3b + c - d)/c = -0.15 \implies \tan^2 \theta_{12} = 0.45, \quad (21)$$

in excellent agreement with data [8].

The limit $\Delta m_{sol}^2 = 0$ implies $2a + b + d = 0$, and

$$\begin{aligned} \Delta m_{atm}^2 &\equiv m_3^2 - (m_2^2 + m_1^2)/2 = (a - b - d)^2 - (d - 3b)^2/4 - 2c^2 \\ &= 6bd - 2(c^2 - d^2) \simeq [6b - 4(c - d)]c, \end{aligned} \quad (22)$$

which can be either positive or negative, corresponding to a normal or inverted ordering of neutrino masses respectively. If $c - d = 0$, then $\tan^2 \theta_{12} < 1/2$ implies an inverted ordering

as in the models of Ref. [16]. It is also possible to set $B = 0$ alone by choosing $f = 0$ in Eq. (5), so that Eq. (20) becomes

$$\tan 2\theta_{12} = \frac{2\sqrt{2}C}{D}, \quad (23)$$

and $\Delta m_{sol}^2 = 0$ implies

$$\Delta m_{atm}^2 \simeq 9(C - D)C^3(m_D^2/\det\mathcal{M}_N)^2. \quad (24)$$

Thus $\tan^2 \theta_{12} < 1/2$ would also imply an inverted ordering of neutrino masses in this case.

The effective neutrino mass m_{ee} measured in neutrinoless double beta decay is simply given by the magnitude of the $\nu_e\nu_e$ entry of $\mathcal{M}_\nu$, i.e. $|a + 2b|$. Using Eqs. (21), (22), and $|\Delta m_{atm}^2| = 2.5 \times 10^{-3} \text{ eV}^2$, it is in the range 0.10 eV for $c - d = 0$ (inverted ordering) and 0.05 eV (normal ordering) for $b = 0$.

Returning to the condition $2a + b + d = 0$ for $\Delta m_{sol}^2 = 0$, under which $\mathcal{M}_\nu$ of Eq. (13) becomes

$$\mathcal{M}_\nu = \begin{pmatrix} a + 2b & c & c \\ c & a - b & -2a - b \\ c & -2a - b & a - b \end{pmatrix}, \quad (25)$$

it should be noted that this form is invariant under the transformation [17]

$$U\mathcal{M}_\nu U^T = \mathcal{M}_\nu, \quad (26)$$

where

$$U = \begin{pmatrix} i2\sqrt{2}/3 & i/3\sqrt{2} & i/3\sqrt{2} \\ i/3\sqrt{2} & 1/2 - i\sqrt{2}/3 & -1/2 - i\sqrt{2}/3 \\ i/3\sqrt{2} & -1/2 - i\sqrt{2}/3 & 1/2 - i\sqrt{2}/3 \end{pmatrix}, \quad U^2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \quad (27)$$

if $c = -2a - 4b$, for which $\tan^2 \theta_{12} = 1/2$. This limit of $\mathcal{M}_\nu$ corresponds thus to a Z_4 symmetry.

In conclusion, the family symmetry S_4 has been advocated as the origin of the observed pattern of neutrino mixing in the context of a complete supersymmetric theory. The key

is the requirement that supersymmetry is unbroken at the seesaw scale where S_4 is broken. This fixes the pattern of S_4 breaking to retain a residual Z_2 (as well as an approximate S_3) symmetry in the Majorana mass matrix of the heavy singlet neutrinos. At the much lower scale of supersymmetry breaking, S_4 as well as Z_2 are allowed to be broken by soft terms. However, the S_4 -invariant Yukawa terms ensure that the charged-lepton mass matrix is diagonal, and the neutrino Dirac masses are equal. This results in a Majorana neutrino mass matrix which exhibits maximal $\nu_\mu - \nu_\tau$ mixing and zero U_{e3} . In the S_3 limit, $(\nu_e + \nu_\mu + \nu_\tau)/\sqrt{3}$ is a mass eigenstate, thus predicting $\tan^2 \theta_{12} = 1/2$, but $\Delta m_{atm}^2 = 0$ as well. Allowing a small deviation from this limit can result in $\tan^2 \theta_{12} = 0.45$ and a nonzero Δm_{atm}^2 , in excellent agreement with data.

I thank Michele Frigerio for an important comment. This work was supported in part by the U. S. Department of Energy under Grant No. DE-FG03-94ER40837.

Appendix The matrices of the $\underline{3}$ representation of S_4 are given by

$$C_1 : \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (28)$$

$$C_2 : \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (29)$$

$$C_3 : \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & -1 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ 1 & 0 & 0 \end{pmatrix}, \\ \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 \\ -1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (30)$$

$$C_4 : \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \\ \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad (31)$$

$$C_5 : \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (32)$$

The matrices of the $\underline{3}'$ representation are the same as those of $\underline{3}$ for $C_{1,2,3}$ and opposite in sign for $C_{4,5}$. Those of the $\underline{2}$ representation are the same as in S_3 , i.e.

$$C_{1,2} : \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (33)$$

$$C_3 : \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}, \begin{pmatrix} \omega^2 & 0 \\ 0 & \omega \end{pmatrix}, \quad (34)$$

$$C_{4,5} : \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \omega \\ \omega^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \omega^2 \\ \omega & 0 \end{pmatrix}, \quad (35)$$

each appearing four times.

References

- [1] Y. Ashie *et al.*, Phys. Rev. **D71**, 112005 (2005) and references therein.
- [2] See for example S.-L. Chen, M. Frigerio, and E. Ma, Phys. Lett. **B612**, 29 (2005) and references therein.
- [3] See for example J. Kubo, A. Mondragon, M. Mondargon, and E. Rodriguez-Jauregui, Prog. Theor. Phys. **109**, 795 (2003); S.-L. Chen, M. Frigerio, and E. Ma, Phys. Rev. **D70**, 073008 (2004) and references therein.
- [4] See for example W. Grimus, A. S. Joshipura, S. Kaneko, L. Lavoura, and M. Tanimoto, JHEP **0407**, 078 (2004).
- [5] See for example E. Ma, hep-ph/0508099 and references therein.
- [6] M. Frigerio, S. Kaneko, E. Ma, and M. Tanimoto, Phys. Rev. **D71**, 011901(R) (2005).
The quaternion group is denoted here as Q_8 , but is called Q_4 in the convention of Ref. [7].
- [7] K. S. Babu and J. Kubo, Phys. Rev. **D71**, 056006 (2005); J. Kubo, hep-ph/0506043.
- [8] B. Aharmim *et al.*, nucl-ex/0502021.
- [9] See for example G. Altarelli and F. Feruglio, Nucl.Phys. **B720**, 64 (2005); K. S. Babu and X.-G. He, hep-ph/0507217.
- [10] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Z. Freedman (North-Holland, Amsterdam, 1979), p. 315; T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK Report No. 79-18, Tsukuba,

- Japan, 1979), p. 95; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [11] Y. Yamanaka, H. Sugawara, and S. Pakvasa, Phys. Rev. **D25**, 1895 (1982); Erratum: *ibid.* **D29**, 2135 (1984).
- [12] E. Ma, Mod. Phys. Lett. **A17**, 2361 (2002).
- [13] E. Ma and G. Rajasekaran, Phys. Rev. **D64**, 113012 (2001).
- [14] E. Ma, Phys. Rev. **D66**, 117301 (2002).
- [15] P. F. Harrison, D. H. Perkins, and W. G. Scott, Phys. Lett. **B530**, 167 (2002).
- [16] S.-L. Chen, M. Frigerio, and E. Ma, hep-ph/0504181, M. Hirsch, A. Villanova del Moral, J. W. F. Valle, and E. Ma, hep-ph/0507148.
- [17] E. Ma, Phys. Rev. Lett. **90**, 221802 (2003); E. Ma and G. Rajasekaran, Phys. Rev. **D68**, 031702(R) (2003).